

DEPARTMENT of ENVIRONMENTAL SERVICES
Water Supply & Pollution Control Division - Biology Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

Lake: BATSON POND	Lake Area (ha):	6.15
Town: WOLFEBORO	Maximum depth (m):	2.7
County: Carroll	Mean depth (m):	1.1
River Basin: Merrimack	Volume (m ³):	68000
Latitude: 43°40'20" N	Relative depth:	1.0
Longitude: 71°09'14" W	Shore configuration:	1.71
Elevation (ft): 998	Areal water load (m/yr):	6.49
Shore length (m): 1500	Flushing rate (yr ⁻¹):	5.90
Watershed area (ha): 69.0	P retention coeff.:	0.61
% watershed ponded: 0.0	Lake type:	natural

<u>BIOLOGICAL:</u>		25 January 1994	19 July 1993
DOM. PHYTOPLANKTON (% TOTAL)	#1	SPARSE - NO DOMINANT	CERATIUM 25%
	#2		SPHAEROCYSTIS 20%
	#3		FILAM. BL-GR SPP 20%
PHYTOPLANKTON ABUNDANCE (cells/mL)			840
CHLOROPHYLL-A (µg/L)			4.88
DOM. ZOOPLANKTON (% TOTAL)	#1	KERATELLA 50%	NAUPLIUS LARVA 56%
	#2	CALANOID COPEPOD 50%	CALANOID COPEPOD 18%
	#3		KERATELLA 13%
ROTIFERS/LITER		6	38
MICROCRUSTACEA/LITER		6	262
ZOOPLANKTON ABUNDANCE (#/L)		12	302
VASCULAR PLANT ABUNDANCE			Common/Abun
SECCHI DISK TRANSPARENCY (m)			2.1
BOTTOM DISSOLVED OXYGEN (mg/L)		5.9	5.8
BACTERIA (E. coli, #/100 ml)	#1		1
	#2		
	#3		

SUMMER THERMAL STRATIFICATION:

not stratified

Depth of thermocline (m): None
Hypolimnion volume (m³) : None
Anoxic volume (m³) : None

CHEMICAL:Lake: BATSON POND
Town: WOLFEBORO

	25 January 1994		19 July 1993		
DEPTH (m)	1.5		1.5		
pH (units)	5.8		6.1		
A.N.C. (Alkalinity)	3.1		1.8		
NITRATE NITROGEN	0.09		< 0.02		
TOTAL KJELDAHL NITROGEN	0.59		0.48		
TOTAL PHOSPHORUS	0.027		0.027		
CONDUCTIVITY (μ mhos/cm)	26.8		18.0		
APPARENT COLOR (cpu)	65		65		
MAGNESIUM			0.34		
CALCIUM			1.1		
SODIUM			1.4		
POTASSIUM			< 0.40		
CHLORIDE	< 2		< 3		
SULFATE	4		3		
TN : TP	25		18		
CALCITE SATURATION INDEX			4.9		

All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 1993

D.O. S.D. PLANT CHL TOTAL CLASS


**	3	4	1	8	Meso.
----	---	---	---	---	-------

COMMENTS:

1. This was an essentially private (privately-owned land surrounded the pond), relatively inaccessible pond that was sampled cooperatively with the NH Fish and Game Department.
2. Pond bottom had a very gradual slope -- no steep-sided drop-offs.
3. Schroederia (60%) was the dominant wholewater phytoplankton genus, with green algae (80%) the dominant class.

Batson Pond

Wolfeboro



0

0.2

Km

[illegible]

TOWN: WOLFEBORO
WEATHER: CLOUDY, WARM & CALM


[illegible]

COMMENTS :

***Dissolved oxygen values are in mg/L**

Batson Pond

Wolfeboro



0 0.2
Km

[illegible]